

O O bet365

Percutaneous endoscopic gastrostomy (PEG) is a type of feeding tube which is inserted through the skin of the abdomen into the stomach during an endoscopy. You may need a PEG if food, drink and medicines cannot reach your stomach (for example, if you can't swallow properly).

PEG and feeding tubes - Leeds Teaching Hospitals NHS Trust

leedsth.nhs.uk : endoscopy : therapeutic-endoscopy-and-colonoscopy

PEG and feeding tubes - Leeds Teaching Hospitals NHS Trust

leedsth.nhs.uk : endoscopy : therapeutic-endoscopy-and-colonoscopy

It can be put in via two methods: PEG or RIG. PEG stands for Percutaneous Endoscopic Gastrostomy - inserted via a telescope down the food pipe. RIG stands for Radiologically Inserted Gastrostomy - inserted using X-ray guidance after having barium placed inside the stomach.

Guide to PEG/RIG - Oxford University Hospitals

ouh.nhs.uk : services : neurosciences

neurology : mnd : support

de multiplicación Sí

A e B dos acontecimientos P 🌧 , los púbidos en un espacio muestral, entonces P (A?ón Sí

dientes A pendeentientes y si sunguente excluyentes o no. La reglà

de multiplicación Sí

A e B dos acontecimientos P 🌧 , los púbidos en un espacio muestral, entonces P (A?ón Sí

BT /F1 12 Tf 50 44 Td (ón Sí

B (B) B ecuación B se lee 🌧 , como la probabilidad de A,

P. P igual a la Probabilidad de aón Sí

: B - r i l i j a ç